

USAID Country Health Statistical Report

Cambodia

May 2008



Table of Contents

1. About the Report

2. Country Profile

I. Statistical Overview

- Demographic Indicators
- Socioeconomic Indicators
- Family Planning Indicators
- Maternal Health Indicators
- Child Survival Indicators
- HIV/AIDS Prevention Indicators
- Infectious Diseases Control Indicators

II. Trends in Health Indicators

- Current and Projected Population by Age and Sex (Figure 1 and Table 1.1)
- Population Estimates and Projections (Table 1.2)
- Estimated Total Fertility Rate per Woman (Figure 2 and Table 2)
- Infant Mortality Rates / Under-5 Mortality Rates (Figure 3 and Table 3)
- Vaccination Coverage (DPT3) Trends (Figure 4 and Table 4)
- Contraceptive Prevalence Rates Married Women (Figure 5 and Table 5)

3. Data Notes

4. Sources

About the Report

Released May 2008

This report is one of a series of Country Health Statistical Reports produced on behalf of the United States Agency for International

Development (USAID) by the Analysis, Information Management and Communication Activity (AIM) Project. Each profile contains

statistical data on current health conditions, population dynamics, health and family planning behavior, and health and population

trends in a given developing country. Information is compiled from AIM's health statistics database, which draws data from a diverse

range of sources listed at the end of this profile.

Hard copy editions of USAID's profiles are available from AIM'S publications department. Reports are available on the USAID Global

Health Web site in Portable Document Format (pdf). Any feedback you have on the content or presentation of this report would be

greatly appreciated. We would also appreciate receiving any more recent, more accurate, or more representative information. Contact

us at the following:

Analysis, Information Management

&

Communications Activity (AIM)

655 15th Street, NW, Suite 450 Washington, DC 20005

Phone: 202-661-8600

Fax: 202-661-8601

E-mail: dolphn@aimglobalhealth.org

The Analysis, Information Management & Communications (AIM) Activity, a USAID contract managed by MasiMax Resources, Inc.

with John Snow, Inc., ORC Macro, and Insight Systems Corporation prepared this document under task order number GPO-M-00-05-

00043-00, with the Bureau for Global Health, USAID.

I. Statistical Overview

*see data notes

Demographic Indicators	Demographic Indicators								
Indicator	Value	Data Unit	Year	Source					
Total Population	14,241,640		2008	BUCEN-IDB-2008					
Population Growth Rate	1.8	%	2008	BUCEN-IDB-2008					
Percent Urban	19	%	2004	World Bank/WDI-2006					
Women, 15-19	905,079		2008	BUCEN-IDB-2008					
Women, 15-49	3,964,797		2008	BUCEN-IDB-2008					
Life Expectancy at Birth	61.7		2008	BUCEN-IDB-2008					
Crude Birth Rate	25.7	per 1,000	2008	BUCEN-IDB-2008					
Crude Death Rate	8.2	per 1,000	2008	BUCEN-IDB-2008					
Number of Live Births	365,725		2008	BUCEN-IDB-2008					
Healthy Life Expectancy: Female	49.5		2002	WHO World Health Report-2004					
Healthy Life Expectancy: Male	45.6		2002	WHO World Health Report-2004					
Population: Percent < Age 15	33.2	%	2008	BUCEN-IDB-2008					

Socioeconomic Indicators				
Indicator	Value	Data Unit	Year	Source
GNI per Capita (PPP)	2,490 1	\$	2005	World Bank/WDI-2006
Health Expenditure as Percentage of GDP	10.9	%	2003	World Bank/WDI-2006
Physicians per 1,000 People	0.2	per 1,000	1997-2004	World Bank/WDI-2006
Adult Literacy Rate	73.6	%	2004	World Bank/WDI Database-2007
Adult Literacy Rate, Female	64 ²	%	2002	World Bank/WDI-2006
Adult Literacy Rate, Male	85 ³	%	2002	World Bank/WDI-2006
Gross Enrollment Ratio - Primary School	134.1	%	2005	UNESCO EFA Global Monitoring Report- 2008
Gender Parity Index - Net Enrollment Ratio	0.98		2005	UNESCO EFA Global Monitoring Report- 2008
Access to an Improved Water Source (Rural)	29	%	2002	World Bank/WDI-2005
Access to an Improved Water Source (Urban)	58	%	2002	World Bank/WDI-2005
Access to Improved Sanitation Facilities (Rural)	8	%	2002	World Bank/WDI-2005
Access to Improved Sanitation Facilities (Urban)	53	%	2002	World Bank/WDI-2005
Population Living Below \$1 a Day	34.1	%	1990-2005	Human Development Report - 2007/2008- 2008
Human Development Index	0.6		2005	Human Development Report - 2007/2008- 2008
Population Below Poverty Line	35	%	2004	CIA World Factbook-2008
Real GDP (Growth) Rate	9.1 ⁴	%	2007	CIA World Factbook-2008
Income Inequality (Gini index)	41.7 ⁵		2004	CIA World Factbook-2008
Unemployment Rate	2.5 ⁶	%	2000	CIA World Factbook-2008
Inflation Rate	4.4 ⁷	%	2007	CIA World Factbook-2008
Airports	17		2007	CIA World Factbook-2008
Airports with paved runways	6		2007	CIA World Factbook-2008

¹ Estimate is based on regression; other PPP figures are extrapolated from the latest International Comparison Programme benchmark estimates.

² Based on survey data.

³ Based on survey data ⁴ estimate

⁵ Estimate

⁶ estimate

⁷ Estimate

Family Planning Indicators				
Indicator	Value	Data Unit	Year	Source
Total Fertility Rate (BUCEN)	3.1		2008	BUCEN-IDB-2008
Total Fertility Rate (DHS)	3.4		2005	Cambodia DHS-2005
Contraceptive Prevalence Rate, Modern Methods, All Women	16.4	%	2005	Cambodia DHS-2005
Contraceptive Prevalence Rate, Modern Methods, Married Women	27.2	%	2005	Cambodia DHS-2005
Median Age of Sexual Debut Among Women, Ages 25-49	20.4		2005	Cambodia DHS-2005
Mean Ideal Family Size	3.3		2005	Cambodia DHS-2005
Women 20-24 Who Gave Birth Before Age 20	26.3	%	2005	Cambodia DHS-2005
Total Fertility Rate, Rural	3.5		2005	DHS STATcompiler as of March 2008-2008
Unmet Need for Family Planning: Rural	25.7	%	2005	DHS STATcompiler as of March 2008-2008
Unmet Need for Family Planning: Urban	21.8	%	2005	DHS STATcompiler as of March 2008-2008
Contraceptive Prevalence Rate, All Methods	40	%	2005	DHS STATcompiler as of March 2008-2008
Equity in use of modern contraception	0.68	Ratio	2005	Cambodia DHS-2005
Percent of need satisfied by modern methods of family planning	41.8	%	2005	Cambodia DHS-2005

Maternal Health Indicators				
Indicator	Value	Data Unit	Year	Source
Maternal Mortality Ratio (WHO/Hill)	540	Per 100,000 live births	2005	WHO/Hill-2005
Maternal Mortality Ratio (DHS)	472 ¹	Per 100,000 live births	2005	Cambodia DHS-2005
Antenatal Care (at least 1 visit)	71.4 ²	%	2005	Cambodia DHS-2005
Antenatal Care (2+ visits)	60.2 ³	%	2005	Cambodia DHS-2005
Antenatal Care (4+ visits)	27 4	%	2005	Cambodia DHS-2005
Assisted Delivery by a Health Professional	43.8	%	2005	Cambodia DHS-2005
Assisted Delivery by Doctor	6.5	%	2005	Cambodia DHS-2005
Assisted Delivery by Other Health Professional	37.3	%	2005	Cambodia DHS-2005
Anemia prevalence among women of reproductive age	46.6	%	2005	Cambodia DHS-2005
Equity in skilled attendance at delivery	0.23	Ratio	2005	Cambodia DHS-2005

 ¹ The value for this country represents data from the 0-6 year period preceding the survey.
 ² The value for this country represents data from the five-year period preceding the survey.
 ³ The value for this country represents data from the five-year period preceding the survey.
 ⁴ The value for this country represents data from the five-year period preceding the survey.

Child Survival Indicators				
Indicator	Value	Data Unit	Year	Source
Acute Respiratory Infection (ARI) and Oral Rehydra	ition The	rapy (ORT)		
ARI Care Seeking - Children Under 5	48.3	%	2005	Cambodia DHS-2005
ORT Use Rate (ORS, RHS, or Increased Fluids)	58.4	%	2005	Cambodia DHS-2005
Mortality Indicators				
Infant Mortality Rate (BUCEN)	56.6	per 1,000 live births	2008	BUCEN-IDB-2008
Infant Mortality Rate (UNICEF)	65		2006	UNICEF-2008
Infant Mortality Rate (DHS)	66	per 1,000 live births	2005	Cambodia DHS-2005
Infant Mortality Rate, Females (BUCEN)	49.1	per 1,000 live births	2008	BUCEN-IDB-2008
Infant Mortality Rate, Males (BUCEN)	63.8	per 1,000 live births	2008	BUCEN-IDB-2008
Under-5 Mortality Rate (BUCEN)	89.1	per 1,000 live births	2006	BUCEN-IDB-2006
Neonatal Mortality Rate	28	per 1000	2005	Cambodia DHS - Preliminary Report- 2005
Under-5 Mortality Rate (UNICEF)	82	per 1,000 live births	2006	UNICEF-2008
Under-5 Mortality Rate (DHS)	83	per 1,000 live births	2005	Cambodia DHS-2005
Nutrition Indicators				
Exclusive Breastfeeding (under 4 mos.)	67.7	%	2005	Cambodia DHS-2005
Exclusive Breastfeeding (under 6 mos.)	60	%	2005	Cambodia DHS-2005
Stunted (height-for-age)	37.3	%	2005	Cambodia DHS-2005
Underweight (weight-for-age)	35.6	%	2005	Cambodia DHS-2005
Wasted (weight-for-height)	7.3	%	2005	Cambodia DHS-2005
Vaccination Coverage				
DPT3 Vaccination Rate (DHS)	78.3	%	2005	Cambodia DHS-2005
DPT3 Vaccination Rate (WHO)	80	%	2006	WHO/Global Summary-2008
Measles Vaccination Rate (DHS)	76.9	%	2005	Cambodia DHS-2005
Measles Vaccination Rate (WHO)	78	%	2006	WHO/Global Summary-2008
Polio Vaccination Rate (DHS)	76.9	%	2005	Cambodia DHS-2005
Polio Vaccination Rate (WHO)	80	%	2006	WHO/Global Summary-2008
Tetanus Toxoid Vaccination	54.2	%	2005	Cambodia DHS-2005
Equity in DPT3 coverage (DHS)	0.78	Ratio	2005	Cambodia DHS-2005
Improved sanitation/hygiene practices	57.9 ¹	%	2005	Cambodia DHS-2005
Vitamin A supplementation coverage among children 6-59 months	34.5	%	2005	Cambodia DHS-2005

¹ The value for this country represents data more likely to be contained by child used, put, or rinsed into toilet or latrine and buried.

HIV/AIDS Prevention Indicators									
Indicator	Value	Data Unit	Year	Source					
HIV Prevalence proportion: Adults (15–49 years)	1.6	%	2005	UNAIDS 2006-2006					
Estimated number of people living with HIV: Adults and Children	130,000		2005	UNAIDS 2006-2006					
Estimated number of people living with HIV: Women (15+ years)	59,000		2005	UNAIDS 2006-2006					
Estimated number of people living with HIV: Children (0–14 years)	7,300		2003	UNAIDS-2004					
Males Reporting Condom Use With Last Non- Regular Partner	82	%	2005	Cambodia DHS - Preliminary Report-2005					
Females Reporting Condom Use With Last Non-Regular Partner	Data Not Available								

Infectious Diseases Control Indicators						
Indicator	Value	Data Unit	Year	Source		
TB Estimated Number of Cases	70,949		2006	WHO Global Health Atlas-2008		
TB Case Detection Rate	62	%	2006	WHO Global Health Atlas-2008		
TB (DOTS) Treatment Success Rate	93	%	2005	WHO Global Health Atlas-2008		
Malaria Cases per 100,000	506 ¹	per 100,000	2003	WHO Global Health Atlas-2008		
DOTS Coverage	100	%	2006	WHO Global Health Atlas-2008		
HIV-Infected with Active TB	9.6	%	2006	WHO Global Health Atlas-2008		
Avian Influenza: Cumulative Number of Confirmed Human Cases	<0.01	Reported # of cases	2008	Epidemic and Pandemic Alert and Response (EPR) section of the WHO website-2008		
Avian Influenza: Cumulative Number of Confirmed Human Deaths	<0.01	Reported # of cases	2008	Epidemic and Pandemic Alert and Response (EPR) section of the WHO website-2008		
Child Sleeps Under Insecticide-Treated Net	4.2	%	2005	DHS STATcompiler as of March 2008-2008		
Equity in Bednet (ITN) Ownership	Data Not Available					

Calculated using (100,000/WHO population)xMalaria Reported # of Cases




Figure 1: Current and Projected Population by Age and Sex (in thousands)

Table 1.1: Current and Projected Population by Age and Sex

Source: BUCEN-IDB Age Group Male - 2000 Male - 2020 Female - 2000 Female - 2020 0-4 784,561 1,006,943 766,854 976,647 5-9 905,289 948,005 887,869 923,527 10-14 883,573 841,203 867,870 822,078 15-19 768,562 770,322 756,271 755,207 20-24 454,152 750,372 447,760 740,766 25-29 461,622 870,507 463,870 859,895 30-34 454,045 836.124 463,167 822,176 35-39 388,970 709,386 407,995 694,552 40-44 220,184 408,946 318,818 398,688 45-49 181,470 400,248 265,704 402,953 50-54 144,802 379,622 201,509 401,738 55-59 111,948 310,604 158,737 351,814 60-64 89,911 163,993 131,801 266,715 65-69 66,894 120,687 103,836 207,480 70-74 48,851 81,015 76,586 138,447 75-79 30,264 47,008 43,878 87,006 80+ 15,573 33,627 22,384 72,499

TABLE 1.2: Population Estimates and Projections

Source: BUCEN-IDB

1960	1970	1980	1990	2000	2010	2020	2030	2040	2050
5,761,223	7,395,691	6,888,021	9,344,566	12,395,580	14,753,320	17,600,800	20,182,594	22,278,723	23,965,562

Figure 2: Estimated Total Fertility Rate per Woman




Table 2: Estimated Total Fertility Rate per Woman

Source: BUCEN-IDB

1970 1980 1990 2000 2010 2020 2030 2040 2050

6.5 7.4 6.0 3.7 3.0 2.7 2.4 2.1 1.8

Figure 3: Infant Mortality Rates / Under-5 Mortality Rates




Table 3: Infant Mortality Rates / Under-5 Mortality Rates

Source: BUCEN-IDB 2008

Indicator	1970	1980	1990	2000	2010	2020	2030	2040	2050
Infant Mortality Rate (BUCEN)	198.9	227.8	143.5	82.2	53.0	37.8	26.6	18.8	13.5
Under-5 Mortality Rate (BUCEN)	264.2	309.4	155.3	110.1	80.2	58.0	40.3	27.2	18.1

Figure 4: Vaccination Coverage (DPT3) Trends




Table 4: Vaccination Coverage (DPT3) Trends

Source: WHO/Global Summary

1985	1990	1995	2000	2005	2006
23.0	38.0	39.0	59.0	82.0	80.0

Figure 5: Contraceptive Prevalence Rates, Modern Methods, Married Women




Table 5: Contraceptive Prevalence Rates, Modern Methods, Married Women

Source: Cambodia DHS

2000	2005
18.8	27.2

Data Notes

*Indicator definitions that differ from those in the data notes section are identified by footnotes accompanying the indicator tables.

Demographic Indicators	Den	noara	phic	Indica	itors
------------------------	-----	-------	------	--------	-------

Total Population The number of people in a given area (i.e., country) in a particular time period (usually a midyear

estimate).

Population Growth Rate The average annual growth rate is the rate of natural increase in a population plus the net migration

rate. The rate of natural increase is the difference between the birth rate and the death rate, but it is

conventionally measured in percentage terms (per hundred rather than per thousand).

Percent Urban The percentage of the midyear population living in areas defined as urban in each country and reported

to the United Nations.

Women, 15-19 The number of women between the ages of 15-19 in the midyear population.

Women, 15-49 The number of women between the ages of 15-49 in the midyear population.

Life Expectancy at Birth The average number of years that a person at age 0 will live if age-specific death rates remain constant.

Life expectancy at birth is highly affected by rates of infant and child death.

Crude Birth Rate The number of births per thousand of the population. The product of the number of live births, divided by

the midpoint population, multiplied by 1,000.

Crude Death Rate The number of deaths per thousand of the population. The product of the number of deaths divided by

the midpoint population, multiplied by 1,000.

Number of Live Births The number of live births, annually, within a country.

Healthy Life Expectancy:

Female

The number of years of in full health that a female newborn can expect to live based on current rates of ill-health and mortality. HALE is based on life expectancy at birth but includes an adjustment for time spent in poor health.

Healthy Life Expectancy:

Male

The number of years of in full health that a male newborn can expect to live based on current rates of ill-health and mortality. HALE is based on life expectancy at birth but includes an adjustment for time spent

in poor health.

Population: Percent < Age 15

Percent of population less than 15 years of age.

Socioeconomic Indicators

GNI per Capita (PPP) The gross national income (GNI) converted to international dollars using purchasing power parity (PPP)

rates. An international dollar has the same purchasing power over GNI as a U.S. dollar has in the

United States.

Health Expenditure as Percentage of GDP

Total health expenditures is the sum of public and private health expenditures figured as a percentage of a country's gross domestic product (GDP). It covers the provision of health services (preventive and curative), family planning activities, nutrition activities, and emergency aid designated for health but

does not include provision of water and sanitation.

Physicians per 1,000

People

Physicians are defined as graduates of any faculty or school of medicine who are working in the country

in any medical field (practice, teaching, research).

Adult Literacy Rate The percentage of people ages 15 and over who can, with understanding, read and write a short, simple

statement about their everyday life.

Adult Literacy Rate,

Female

The percentage of women ages 15 and over who can, with understanding, read and write a short,

simple statement about their everyday life.

Adult Literacy Rate, Male

The percentage of men ages 15 and over who can, with understanding, read and write a short, simple statement about their everyday life.

Gross Enrollment Ratio - Primary School

Total enrolment in primary level of education, regardless of age, expressed as a percentage of the population in the official age group corresponding to primary level of education. The GER can exceed 100% due to late entry or/and repetition.

Gender Parity Index - Net Enrollment Ratio

The ratio of the female-to-male values (or male to female, in certain cases) of net primary school enrollment rates (NER). NER measures the number of pupils in the official age group for a given level of ecudation, expressed as a percentage of the population in that age group.

Access to an Improved Water Source (Rural)

Access to an improved water source in rural areas. Refers to the percentage of the rural population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs.

Access to an Improved Water Source (Urban)

Access to an improved water source in urban areas. Refers to the percentage of the urban population with reasonable access to an adequate amount of water from an improved source, such as a household connection, public standpipe, borehole, protected well or spring, or rainwater collection. Unimproved sources include vendors, tanker trucks, and unprotected wells and springs.

Access to Improved Sanitation Facilities (Rural)

Access to improved sanitation facilities in rural areas. Refers to the percentage of the rural population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.

Access to Improved Sanitation Facilities (Urban)

Access to improved sanitation facilities in urban areas. Refers to the percentage of the urban population with access to at least adequate excreta disposal facilities (private or shared, but not public) that can effectively prevent human, animal, and insect contact with excreta.

Population Living Below \$1 a Day

The percentage of the population living below the specified poverty line of \$1 a day at 1985 international prices (equivalent to \$1.08 at 1993 international prices), adjusted for purchasing power parity.

Human Development Index

The HDI is a composite index that measures the average achievements in a country in three basic dimensions of human development: a long and healthy life, as measured by life expectancy at birth; knowledge, as measured by the adult literacy rate and the combined gross enrollment ratio for primary, secondary, and tertiary schools; and a decent standard of living, as measured by gross domestic product (GDP) per capita in purchasing power parity (PPP) US dollars.

Population Below Poverty Line

National estimates of the percentage of the population falling below the poverty line are based on surveys of sub-groups, with the results weighted by the number of people in each group. Definitions of poverty vary considerably among nations. For example, rich nations generally employ more generous standards of poverty than poor nations.

Real GDP (Growth) Rate

This entry gives the gross domestic product (GDP) growth on an annual basis adjusted for inflation and expressed as a percent.

Income Inequality (Gini index)

This index measures the degree of inequality in the distribution of family income in a country. It is calculated from the Lorenz curve, in which cumulative family income is plotted against the number of families arranged from the poorest to the richest. The index is the ratio of (a) the area between a country's Lorenz curve and the 45 degree helping line to (b) the entire triangular area under the 45 degree line. The more nearly equal a country's income distribution, the closer its Lorenz curve to the 45 degree line and the lower its Gini index, e.g., a Scandinavian country with an index of 25. The more unequal a country's income distribution, the higher its Gini index, e.g., a Sub-Saharan country with an index of 50. If income were distributed with perfect equality, the index would be zero; if income were distributed with perfect inequality, the index would be 100.

Unemployment Rate

This entry contains the percent of the labor force that is without jobs. Substantial underemployment might be noted.

Inflation Rate

This entry furnishes the annual percent change in consumer prices compared with the previous year's consumer prices.

Airports

This entry gives the total number of airports or airfields recognizable from the air. The runway(s) may be paved (concrete or asphalt surfaces) or unpaved (grass, earth, sand, or gravel surfaces) but may include closed or abandoned installations. Airports or airfields that are no longer recognizable (overgrown, no facilities, etc.) are not included. Note that not all airports have accommodations for refueling, maintenance, or air traffic control.

Airports with paved runways

This entry gives the total number of airports with paved runways (concrete or asphalt surfaces) by length. For airports with more than one runway, only the longest runway is included according to the following five groups - (1) over 3,047 m, (2) 2,438 to 3,047 m, (3) 1,524 to 2,437 m, (4) 914 to 1,523 m, and (5) under 914 m. Only airports with usable runways are included in this listing. Not all airports have facilities for refueling, maintenance, or air traffic control.

Family Planning Indicators

Total Fertility Rate (BUCEN)

The number of children a woman between ages 15-49 would have during her reproductive life, if, for all of her childbearing years she were to experience the age-specific birth rates for that given year.

Total Fertility Rate (DHS)

The number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.

Contraceptive Prevalence Rate, Modern Methods, All Women Percentage of all women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).

Contraceptive Prevalence Rate, Modern Methods, Married Women Percent of currently married women ages 15-49 currently using a modern method of contraception. Modern methods include oral contraceptives, IUDs, injectables, female and male sterilization, all emergency contraception, and barrier methods (diaphragm, foam, jelly, male and female condom).

Median Age of Sexual Debut Among Women, Ages 25-49 Median age of first sexual intercourse for women ages 25-49.

Mean Ideal Family Size

Mean ideal number of children for all women, according to number of living children.

Women 20-24 Who Gave Birth Before Age 20

Percentage of women ages 20-24 who have given birth before age 20.

Total Fertility Rate, Rural

In rural areas only, the number of children a woman between ages 15-49 would have during her lifetime if she were to bear children at the currently observed rates.

Unmet Need for Family Planning: Rural

In rural areas, percentage of married women ages 15-49 with unmet need for family planning. Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrheic women who are not using family planning and whose last birth was mistimed, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they want to wait 2 or more years for their next birth. Also included are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth unless they say it would not be a problem if they discovered they were pregnant in the next few weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic women whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and want no more children.

Unmet Need for Family Planning: Urban

In urban areas, percentage of married women ages 15-49 with unmet need for family planning. Unmet need for spacing includes pregnant women whose pregnancy was mistimed, amenorrheic women who are not using family planning and whose last birth was mistimed, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and say they want to wait 2 or more years for their next birth. Also included are fecund women who are not using any method of family planning and say they are unsure whether they want another child or who want another child but are unsure when to have the birth unless they say it would not be a problem if they discovered they were pregnant in the next few weeks. Unmet need for limiting refers to pregnant women whose pregnancy was unwanted, amenorrheic women whose last child was unwanted, and fecund women who are neither pregnant nor amenorrheic and who are not using any method of family planning and want no more children.

Contraceptive Prevalence Rate, All Methods

Equity in use of modern contraception

Contraceptive prevalence rate in the lowest wealth quintile/contraceptive prevalence rate in the highest wealth quintile.

Percentage of currently married women ages 15-49 currently using ANY method of contraception.

Percent of need satisfied by modern methods of family planning

Number of women in union who are fecund and are currently using modern contraceptive methods to stop or postpone the next pregnancy/number of women in union who are fecund and who desire to either terminate childbearing or to postpone their next birth for 2 years or more.

Maternal Health Indicators

Maternal Mortality Ratio (WHO/Hill)

The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births arrived through Hill et al., adjustment procedure depending on the nature of data used. The margins of uncertainty associated with the estimated maternal mortality ratios are very large and the estimates should not be used to monitor trends in the short term (including comparisons between 1995 and 2000 estimates). In addition, cross-country comparisons should be treated with considerable circumspection because different strategies are used to derive the estimates for different countries rendering comparisons fraught with difficulty. For further information please refer to the source.

Maternal Mortality Ratio (DHS)

The estimated number of women who die as a result of pregnancy or childbirth per 100,000 live births, arrived mostly through the "sisterhood method". The data are aggregated based on different time periods ranging from four to ten years preceding the survey. Thus, the data may not be suitable for trend analysis due to over-lapping year ranges of estimates. For further information on methodology please refer to: Stanton, Cynthia, Noureddine Abderrahim, and Kenneth Hill. 1997. DHS Maternal Mortality Indicators: An assessment of Data Quality and Implications for Data Use. Calverton: Macro International Inc., or individual country DHS reports.

Antenatal Care (at least 1 visit)

Percentage of women of reproductive age (15-49) who receive at least one antenatal care visit during pregnancy (in the three-year period preceding the survey).

Antenatal Care (2+ visits)

Percentage of women of reproductive age (15-49) who receive at least two antenatal care visits during pregnancy (in the three-year period preceding the survey).

Antenatal Care (4+ visits)

Percentage of women of reproductive age (15-49) who receive four or more antenatal care visits during pregnancy (in the three-year period preceding the survey).

Assisted Delivery by a Health Professional

The percentage of births/deliveries that occur with the assistance of any trained health professional during the five-year period preceding the survey. May include doctors, nurses, or midwives.

Assisted Delivery by Doctor

The percentage of live births/deliveries that occur with the assistance of a doctor during the five-year period preceding the survey.

Assisted Delivery by Other Health Professional The percentage of live births/deliveries that occur with the assistance of a nurse or midwife during the five-year period preceding the survey.

Anemia prevalence among women of reproductive age

Number of women of reproductive age (15-49 years) with anemia/number of women of reproductive age (15-49 years).

Equity in skilled attendance at delivery

Percentage of births in the lowest wealth quintile attended by a doctor, nurse, or trained midwife/percentage of births in the highest wealth quintile attended by a doctor, nurse, or trained midwife.

Child Survival Indicators

ARI Care Seeking - Children Under 5

Percentage of children under five years who were ill with an acute respiratory infection (ARI), which is associated with cough, rapid breathing and a high fever, during the two weeks preceding the survey, for whom care was sought from a health facility or provider.

ORT Use Rate (ORS, RHS, or Increased Fluids)

Oral Rehydration Therapy (ORT). Percentage of children under age 5 with diarrhea two weeks prior to survey who received increased fluids, oral rehydration solution (ORS) or recommended home solution (RHS).

Infant Mortality Rate (BUCEN)

The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year.

Infant Mortality Rate (UNICEF)	The under five mortality rate is the probability (as expressed as a rate per 1,000 live births)of a child born in a specified year dying before reaching the age of one if subject to current age-specific mortality rates.
Infant Mortality Rate (DHS)	The estimated annual number of deaths of infants under 12 months in a given year per 1,000 live births in that same year (five-year period preceding survey).
Infant Mortality Rate, Females (BUCEN)	The estimated annual number of deaths of female infants under 12 months in a given year per 1,000 live births in that same year.
Infant Mortality Rate, Males (BUCEN)	The estimated annual number of male infants under 12 months in a given year per 1,000 live births in that same year.
Under-5 Mortality Rate (BUCEN)	Annual number of deaths that occur in children 0-4 years old per 1,000 births.
Neonatal Mortality Rate	The estimated number of infant deaths in the first month of life per 1,000 live births in the five-year period preceding the survey.
Under-5 Mortality Rate (UNICEF)	Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).
Under-5 Mortality Rate (DHS)	Annual number of deaths that occur in children 0-4 years old per 1,000 births (five-year period preceding survey).
Exclusive Breastfeeding (under 4 mos.)	Percentage of children under 4 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.
Exclusive Breastfeeding (under 6 mos.)	Percentage of children under 6 months exclusively breastfed. Exclusive breastfeeding is defined as providing no food or liquid other than breast milk to the child during the 24-hour period before the survey.
Stunted (height-for-age)	Percentage of children under age five whose height-for-age is below minus-two standard deviations from the median of the reference population.
Underweight (weight-forage)	Percentage of children under age five whose weight-for-age is below minus-two standard deviations from the median of the reference population.
Wasted (weight-for- height)	Percentage of children under age five whose weight-for-height is below minus-two standard deviations from the median of the reference population.
DPT3 Vaccination Rate (DHS)	Proportion of living children 12-23 months old who have received three complete doses of vaccines against diphtheria, pertussis, and tetanus.
DPT3 Vaccination Rate (WHO)	Proportion of living children 12-23 months old who have received three complete doses of vaccines aginst diphtheria, pertussis, and tetanus.
Measles Vaccination Rate (DHS)	Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.
Measles Vaccination Rate (WHO)	Percentage of living children 12-23 months old who have received one dose of Measles Containing Vaccince.
Polio Vaccination Rate (DHS)	Proportion of living children 12-23 months who have received three doses of polio vaccine.
Polio Vaccination Rate (WHO)	Proportion of living children 12-23 months who have received three doses of polio vaccine.
Tetanus Toxoid Vaccination	Percentage of pregnant women receiving two (or more) doses of tetanus toxoid (three-year period preceding survey).
Equity in DPT3 coverage (DHS)	DPT3 coverage in the lowest wealth quintile/DPT3 coverage in the highest wealth quintile.
Improved sanitation/ hygiene practices	Number of households with child under 5 years of age whose youngest child's feces was safely disposed of the last time he/she passed stool/total number of households with a child under 5 years of age.

Vitamin A supplementation coverage among children 6-59 months Number of children aged 6-59 months who received a vitamin A supplement in the last 6 months/total number of children aged 6-59 months surveyed.

HIV/AIDS Prevention Indicators

HIV Prevalence proportion: Adults (15–49 years)

The adult HIV prevalence proportion is the estimated number of adults living with HIV divided by the adult population (aged 15–49).

Estimated number of people living with HIV: Adults and Children

Estimated number of adults and children living with HIV. Adults are 15 years and over. Children are defined as those aged 0–14 years.

Estimated number of people living with HIV: Women (15+ years)

Estimated number of women (aged 15 and over) living with HIV.

Estimated number of people living with HIV: Children (0–14 years)

Estimated number of children under age 15 living with HIV.

Males Reporting Condom Use With Last Non-Regular Partner Among men who have had high risk sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.

Females Reporting Condom Use With Last Non-Regular Partner Among women who have had high risk sex in the last year, percentage who used condoms during their last sexual intercourse with a non-regular partner/non-cohabitating partner.

Infectious Diseases Control Indicators

TB Estimated Number of Cases

The estimated number of cases (all forms).

TB Case Detection Rate

TB Case Detection Rate is defined as the percentage of the annual new smear-positive notifications of the estimated annual new smear-positive incidence.

TB (DOTS) Treatment Success Rate

The proportion of smear-positive patients who were cured plus the proportion who completed treatment.

Malaria Cases per 100,000

Number of malaria cases per 100,000 of the population during the year for which data was reported.

DOTS Coverage

The percentage of the national population living in areas where health services have adopted DOTS. "Areas" are the lowest administrative or management units in the country (townships, districts, counties, etc). If an area is considered a DOTS area, then all cases registered and reported in that area are considered DOTS cases and the population living within the boundaries of that area counts toward national DOTS coverage. As a measure of patient access to diagnosis and treatment under DOTS, coverage is an approximation, and usually an overestimate.

HIV-Infected with Active TB

The TB incidence rate in HIV-infected people 15-49 divided by the incidence rate in HIV-uninfected people 15-49.

Avian Influenza: Cumulative Number of Confirmed Human Cases The cumulative number of confirmed human cases of Avian Influenza A/(H5N1) reported to WHO. The total number of cases includes number of deaths. WHO reports only laboratory-confirmed cases.

Avian Influenza: Cumulative Number of Confirmed Human Deaths

The cumulative number of confirmed human deaths from Avian Influenza A/(H5N1) reported to WHO. WHO reports only laboratory-confirmed cases and deaths.

Child Sleeps Under Insecticide-Treated Net Percentage of children under five years of age who slept under an Insecticde Treated Net (ITN) the night before the survey. An ITN is (1) a permanent net that does not require any treatment, or (2) a pretreated net obtained within the last six months, or (3) a net that has been soaked with insecticide within the past six months.

Equity in Bednet (ITN)
Ownership

Proportion of households in the lowest income quintile with an ITN / proportion of households in the highest income quintile with an ITN.

Statistical Sources For Cambodia

BUCEN-IDB 2006 United States Census Bureau (BUCEN), International Programs Center. International Database. August 2006.

BUCEN-IDB 2008 United States Census Bureau (BUCEN), International Programs Center. International

Database. February 2008.

Cambodia DHS 2000

National Institute of Statistics, Directorate General for Health [Cambodia], and ORC

Macro. 2001. Cambodia Demographic and Health Survey 2000. Phnom Penh, Cambodia and Calverton, Maryland USA: National Institute of Statistics, Directorate General for Health, and ORC Macro.

Cambodia DHS 2005

National Institute of Public Health, National Institute of Statistics [Cambodia] and ORC

Macro. 2006. Cambodia Demographic and Health Survey 2005. Phnom Penh, Cambodia and Calverton, Maryland, USA: National Institute of Public Health, National Institute of

Statistics and ORC Macro.

Cambodia DHS - Preliminary
Report 2005

National Institute of Public Health, Ministry of Health, Phnom Penh, Cambodia. National Institute of Statistics, Ministry of Planning, Phnom Penh, Cambodia. MEASURE DHS,

ORC Macro, Maryland, USA. July 2006

2008 2008

WHO website 2008

CIA World Factbook 2008 CIA World Factbook, 2008 https://cia.gov/cia//publications/factbook/docs/profileguide. html. Last accessed April, 2008.

DHS STATcompiler as of March

DHS STATcompiler as of March 2008 http://www.statcompiler.com/statcompiler/index.

Epidemic and Pandemic Alert and Response (EPR) section of the WHO Site:http://www.who.int/csr/disease/avian_influenza/country/cases table 2008 03 18/en/index.html

Human Development Report
2007/2008 2008

Human Development Report 2007/2008. Fighting climate change: Human solidarity in a divided world. Published for the United Nations Development Programme (UNDP)

UNAIDS 2004 Report on the Global HIV/AIDS Epidemic: 4th Global Report. July 2004. The Joint United Nations Programme on HIV/AIDS, Geneva.

UNAIDS 2006 2006 2006 2006 Report on the Global AIDS Epidemic. Geneva: UNAIDS, 2006.

UNESCO EFA Global Monitoring
Report 2008. Education for All by 2015 - Will we make it? United
Nations Educational, Scientific and Cultural Organization, Paris.

UNICEF 2008

UNICEF. Progress since the World Summit for Children. http://www.childinfo.org/areas/childmortality/u5data.php. Data accessed on web site in March, 2008

WHO Global Health Atlas 2008 Accessed April of 2008

WHO Global Health Atlas 2008 Accessed March of 2008

WHO World Health Report 2004

The World Health Report 2004: Changing history. Geneva, World Health Organization, 2004. Available at: http://www.who.int/whr/2004/en/index.html. Last accessed February, 2007.

WHO/Global Summary 2002	WHO Vaccine-Preventable Diseases Monitoring System. 2002 Global Summary. World Health Organization, Geneva.
WHO/Global Summary 2008	Immunization surveillance, assessment and monitoring section of the WHO website: http://www.who.int/immunization_monitoring/en/globalsummary/timeseries/ tscoveragedtp3.htm
WHO/Global Summary 2008	Immunization surveillance, assessment and monitoring section of the WHO website: http://www.who.int/immunization_monitoring/en/globalsummary/timeseries/ tscoveragemcv.htm
WHO/Global Summary 2008	Immunization surveillance, assessment and monitoring section of the WHO website: http://www.who.int/immunization_monitoring/en/globalsummary/timeseries/tscoveragepol3.htm
WHO/Hill 2005	Maternal Mortality in 2005: Estimates developed by WHO, UNICEF, UNFPA and The World Bank
World Bank/WDI 2005	World Bank Development Indicators 2005, CD-Rom. Washington, DC: The World Bank, 2005.
World Bank/WDI 2006	World Bank Development Indicators 2006. Washington, DC: The World Bank, 2005. Available at: http://devdata.worldbank.org/wdi2006/contents/index2.htm. Last accessed April, 2007.
World Bank/WDI Database 2007	World Bank Development Indicators Database 2007. Washington, DC: The World Bank, 2007. Available at: http://devdata.worldbank.org/data-query/. Last accessed April, 2007.